

Analytical obstructions to the weak approximation of Sobolev mappings into manifolds

Antoine Detaille

Université Claude Bernard Lyon 1 — Institut Camille Jordan
Website: andetaille.github.io

The 11th of June 2025

In this talk

Main goal: new families of counterexamples to the weak approximation property of Sobolev mappings into manifolds, joint work with J. Van Schaftingen (UCLouvain).

On the way:

- Sobolev mappings: definition and motivation;
- the strong density problem: statement and complete answer;
- the weak approximation problem: statement and state of the art.

Sobolev spaces with values into manifolds

Let \mathcal{N} be a smooth compact Riemannian manifold, isometrically embedded in \mathbb{R}^ν .
Let \mathcal{M} be a smooth compact Riemannian manifold of dimension m and $1 \leq p < +\infty$.

Definition

$$W^{1,p}(\mathcal{M}; \mathcal{N}) = \{u \in W^{1,p}(\mathcal{M}; \mathbb{R}^\nu) : u(x) \in \mathcal{N} \text{ for almost every } x \in \mathcal{M}\}$$

The strong density problem

Theorem

The space $C^\infty(\mathcal{M})$ is dense in $W^{1,p}(\mathcal{M})$.

Question

Define

$$H_S^{1,p}(\mathcal{M}; \mathcal{N}) = \{u \in W^{1,p}(\mathcal{M}; \mathcal{N}) : \text{there exists } (u_n)_{n \in \mathbb{N}} \text{ in } C^\infty(\mathcal{M}; \mathcal{N}) \text{ such that } u_n \rightarrow u\}.$$

Does it hold that $H_S^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$?

The strong density problem

Theorem

The space $C^\infty(\mathcal{M})$ is dense in $W^{1,p}(\mathcal{M})$.

Question

Define

$$H_S^{1,p}(\mathcal{M}; \mathcal{N}) = \{u \in W^{1,p}(\mathcal{M}; \mathcal{N}) : \text{there exists } (u_n)_{n \in \mathbb{N}} \text{ in } C^\infty(\mathcal{M}; \mathcal{N}) \text{ such that } u_n \rightarrow u\}.$$

Does it hold that $H_S^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$?

The strong density theorem

Theorem (Bethuel (1991))

Assume that $p < m$. Then, $H_S^{1,p}(\mathbb{B}^m; \mathcal{N}) = W^{1,p}(\mathbb{B}^m; \mathcal{N})$ if and only if $\pi_{[\rho]}(\mathcal{N}) = \{0\}$.

Extensions to $W^{s,p}$: Brezis and Mironescu (2015, $0 < s < 1$); Bousquet, Ponce, and Van Schaftingen (2015, $s = 2, 3, \dots$); D. (2023, $s > 1$ noninteger).

The case where \mathcal{M} is topologically non-trivial was explored by Hang and Lin (2003).
There, *global* obstructions may arise.

The strong density theorem

Theorem (Bethuel (1991))

Assume that $p < m$. Then, $H_S^{1,p}(\mathbb{B}^m; \mathcal{N}) = W^{1,p}(\mathbb{B}^m; \mathcal{N})$ if and only if $\pi_{[\rho]}(\mathcal{N}) = \{0\}$.

Extensions to $W^{s,p}$: Brezis and Mironescu (2015, $0 < s < 1$); Bousquet, Ponce, and Van Schaftingen (2015, $s = 2, 3, \dots$); D. (2023, $s > 1$ noninteger).

The case where \mathcal{M} is topologically non-trivial was explored by Hang and Lin (2003).
There, *global* obstructions may arise.

The strong density theorem

Theorem (Bethuel (1991))

Assume that $p < m$. Then, $H_S^{1,p}(\mathbb{B}^m; \mathcal{N}) = W^{1,p}(\mathbb{B}^m; \mathcal{N})$ if and only if $\pi_{[\rho]}(\mathcal{N}) = \{0\}$.

Extensions to $W^{s,p}$: Brezis and Mironescu (2015, $0 < s < 1$); Bousquet, Ponce, and Van Schaftingen (2015, $s = 2, 3, \dots$); D. (2023, $s > 1$ noninteger).

The case where \mathcal{M} is topologically non-trivial was explored by Hang and Lin (2003). There, *global* obstructions may arise.

Let's be less demanding: weak approximation

We say that $(u_n)_{n \in \mathbb{N}}$ *weakly converges* to u in $W^{1,p}$, and we write $u_n \rightharpoonup u$, whenever $u_n \rightarrow u$ almost everywhere and

$$\sup_{n \in \mathbb{N}} \mathcal{E}^{1,p}(u_n, \mathcal{M}) = \sup_{n \in \mathbb{N}} \int_{\mathcal{M}} |Du_n|^p < +\infty.$$

Define

$$H_W^{1,p}(\mathcal{M}; \mathcal{N}) = \{u \in W^{1,p}(\mathcal{M}; \mathcal{N}) : \text{there exists } (u_n)_{n \in \mathbb{N}} \text{ in } C^\infty(\mathcal{M}; \mathcal{N}) \text{ such that } u_n \rightharpoonup u\}.$$

Question

Does it hold that $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$?

Let's be less demanding: weak approximation

We say that $(u_n)_{n \in \mathbb{N}}$ *weakly converges* to u in $W^{1,p}$, and we write $u_n \rightharpoonup u$, whenever $u_n \rightarrow u$ almost everywhere and

$$\sup_{n \in \mathbb{N}} \mathcal{E}^{1,p}(u_n, \mathcal{M}) = \sup_{n \in \mathbb{N}} \int_{\mathcal{M}} |Du_n|^p < +\infty.$$

Define

$$H_W^{1,p}(\mathcal{M}; \mathcal{N}) = \{u \in W^{1,p}(\mathcal{M}; \mathcal{N}) : \text{there exists } (u_n)_{n \in \mathbb{N}} \text{ in } C^\infty(\mathcal{M}; \mathcal{N}) \text{ such that } u_n \rightharpoonup u\}.$$

Question

Does it hold that $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$?

A topological obstruction: here we go again?

If $p \notin \mathbb{N}$ and $\pi_{[p]}(\mathcal{N}) \neq \{0\}$, then $H_W^{1,p}(\mathcal{M}; \mathcal{N}) \subsetneq W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever $\dim \mathcal{M} > p$.

Theorem (Bethuel (1991))

If $p \notin \mathbb{N}$, then $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = H_S^{1,p}(\mathcal{M}; \mathcal{N})$.

A topological obstruction: here we go again?

If $p \notin \mathbb{N}$ and $\pi_{[p]}(\mathcal{N}) \neq \{0\}$, then $H_W^{1,p}(\mathcal{M}; \mathcal{N}) \subsetneq W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever $\dim \mathcal{M} > p$.

Theorem (Bethuel (1991))

If $p \notin \mathbb{N}$, then $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = H_S^{1,p}(\mathcal{M}; \mathcal{N})$.

A new phenomenon: the case $p \in \mathbb{N}$

Unlike for $2 < p < 3$, we have $\frac{x}{|x|} \in H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$.

More generally, we have:

- $H_S^{1,2}(\mathbb{B}^3; \mathbb{S}^2) \subsetneq H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2) = W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$ (Bethuel (1990));
- $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever \mathcal{N} is $(p-1)$ -connected (Hajłasz (1994));
- $H_W^{1,2}(\mathcal{M}; \mathcal{N}) = W^{1,2}(\mathcal{M}; \mathcal{N})$ for more general \mathcal{N} (Pakzad and Rivière (2003)).

With more work, we should be able to prove that weak approximation always holds for $p \in \mathbb{N}$. No?

A new phenomenon: the case $p \in \mathbb{N}$

Unlike for $2 < p < 3$, we have $\frac{x}{|x|} \in H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$.

More generally, we have:

- $H_S^{1,2}(\mathbb{B}^3; \mathbb{S}^2) \subsetneq H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2) = W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$ (Bethuel (1990));
- $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever \mathcal{N} is $(p-1)$ -connected (Hajlasz (1994));
- $H_W^{1,2}(\mathcal{M}; \mathcal{N}) = W^{1,2}(\mathcal{M}; \mathcal{N})$ for more general \mathcal{N} (Pakzad and Rivière (2003)).

With more work, we should be able to prove that weak approximation always holds for $p \in \mathbb{N}$. No?

A new phenomenon: the case $p \in \mathbb{N}$

Unlike for $2 < p < 3$, we have $\frac{x}{|x|} \in H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$.

More generally, we have:

- $H_S^{1,2}(\mathbb{B}^3; \mathbb{S}^2) \subsetneq H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2) = W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$ (Bethuel (1990));
- $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever \mathcal{N} is $(p-1)$ -connected (Hajłasz (1994));
- $H_W^{1,2}(\mathcal{M}; \mathcal{N}) = W^{1,2}(\mathcal{M}; \mathcal{N})$ for more general \mathcal{N} (Pakzad and Rivière (2003)).

With more work, we should be able to prove that weak approximation always holds for $p \in \mathbb{N}$. No?

A new phenomenon: the case $p \in \mathbb{N}$

Unlike for $2 < p < 3$, we have $\frac{x}{|x|} \in H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$.

More generally, we have:

- $H_S^{1,2}(\mathbb{B}^3; \mathbb{S}^2) \subsetneq H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2) = W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$ (Bethuel (1990));
- $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever \mathcal{N} is $(p-1)$ -connected (Hajłasz (1994));
- $H_W^{1,2}(\mathcal{M}; \mathcal{N}) = W^{1,2}(\mathcal{M}; \mathcal{N})$ for more general \mathcal{N} (Pakzad and Rivière (2003)).

With more work, we should be able to prove that weak approximation always holds for $p \in \mathbb{N}$. No?

A new phenomenon: the case $p \in \mathbb{N}$

Unlike for $2 < p < 3$, we have $\frac{x}{|x|} \in H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$.

More generally, we have:

- $H_S^{1,2}(\mathbb{B}^3; \mathbb{S}^2) \subsetneq H_W^{1,2}(\mathbb{B}^3; \mathbb{S}^2) = W^{1,2}(\mathbb{B}^3; \mathbb{S}^2)$ (Bethuel (1990));
- $H_W^{1,p}(\mathcal{M}; \mathcal{N}) = W^{1,p}(\mathcal{M}; \mathcal{N})$ whenever \mathcal{N} is $(p-1)$ -connected (Hajłasz (1994));
- $H_W^{1,2}(\mathcal{M}; \mathcal{N}) = W^{1,2}(\mathcal{M}; \mathcal{N})$ for more general \mathcal{N} (Pakzad and Rivière (2003)).

With more work, we should be able to prove that weak approximation always holds for $p \in \mathbb{N}$. No? 😞

Obstructions strike back: the analytical obstruction

Theorem (Bethuel (2020))

If $m \geq 4$, then $H_W^{1,3}(\mathcal{M}; \mathbb{S}^2) \subsetneq W^{1,3}(\mathcal{M}; \mathbb{S}^2)$.

Global topological obstructions were already known (Hang and Lin (2003)).

Here, the obstruction is local: it arises already if $\mathcal{M} = \mathbb{B}^4$.

Ingredients involve: the Hopf invariant, Pontryagin construction, the theory of scans by Hardt and Rivière (2003), and branched optimal transportation.

Obstructions strike back: the analytical obstruction

Theorem (Bethuel (2020))

If $m \geq 4$, then $H_W^{1,3}(\mathcal{M}; \mathbb{S}^2) \subsetneq W^{1,3}(\mathcal{M}; \mathbb{S}^2)$.

Global topological obstructions were already known (Hang and Lin (2003)).

Here, the obstruction is local: it arises already if $\mathcal{M} = \mathbb{B}^4$.

Ingredients involve: the Hopf invariant, Pontryagin construction, the theory of scans by Hardt and Rivière (2003), and branched optimal transportation.

Obstructions strike back: the analytical obstruction

Theorem (Bethuel (2020))

If $m \geq 4$, then $H_W^{1,3}(\mathcal{M}; \mathbb{S}^2) \subsetneq W^{1,3}(\mathcal{M}; \mathbb{S}^2)$.

Global topological obstructions were already known (Hang and Lin (2003)).

Here, the obstruction is local: it arises already if $\mathcal{M} = \mathbb{B}^4$.

Ingredients involve: the Hopf invariant, Pontryagin construction, the theory of scans by Hardt and Rivière (2003), and branched optimal transportation.

An analytical obstruction for every $p \in \mathbb{N} \setminus \{0, 1\}$

Theorem (D. and Van Schaftingen (2024))

For every $p \in \mathbb{N} \setminus \{0, 1\}$, there exists a compact Riemannian manifold \mathcal{N} such that, if $\dim \mathcal{M} > p$, then

$$H_W^{1,p}(\mathcal{M}; \mathcal{N}) \subsetneq W^{1,p}(\mathcal{M}; \mathcal{N}).$$

Key procedure in the proof: superlinear energy growth

The *relaxed energy* is defined as

$$\mathcal{E}_{\text{rel}}^{1,p}(u, \mathcal{M}) = \inf \liminf_{n \rightarrow +\infty} \int_{\mathcal{M}} |\mathrm{D}u_n|^p,$$

where the \inf is over all sequences of $C^\infty(\mathcal{M}; \mathcal{N})$ maps converging a.e. to u .

We construct a sequence $(u_n)_{n \in \mathbb{N}}$ such that

$$\liminf_{n \rightarrow +\infty} \frac{\mathcal{E}_{\text{rel}}^{1,p}(u_n, \mathcal{M})}{\mathcal{E}^{1,p}(u_n, \mathcal{M})} = +\infty.$$

The conclusion follows from the nonlinear uniform boundedness principle (Hang and Lin (2003), Monteil and Van Schaftingen (2019)), a nonlinear version of the Banach–Steinhaus theorem.

Key procedure in the proof: superlinear energy growth

The *relaxed energy* is defined as

$$\mathcal{E}_{\text{rel}}^{1,p}(u, \mathcal{M}) = \inf \liminf_{n \rightarrow +\infty} \int_{\mathcal{M}} |\mathrm{D}u_n|^p,$$

where the \inf is over all sequences of $C^\infty(\mathcal{M}; \mathcal{N})$ maps converging a.e. to u .

We construct a sequence $(u_n)_{n \in \mathbb{N}}$ such that

$$\liminf_{n \rightarrow +\infty} \frac{\mathcal{E}_{\text{rel}}^{1,p}(u_n, \mathcal{M})}{\mathcal{E}^{1,p}(u_n, \mathcal{M})} = +\infty.$$

The conclusion follows from the nonlinear uniform boundedness principle (Hang and Lin (2003), Monteil and Van Schaftingen (2019)), a nonlinear version of the Banach–Steinhaus theorem.

A second family of analytical obstructions

Theorem (D. and Van Schaftingen (2024))

For every $n \in \mathbb{N}_*$, if $\dim \mathcal{M} > 4n - 1$, then

$$H_W^{1,4n-1}(\mathcal{M}; \mathbb{S}^{2n}) \subsetneq W^{1,4n-1}(\mathcal{M}; \mathbb{S}^{2n}).$$

The key ingredient is a periodic construction using a Whitehead product.

This shows that Bethuel's counterexample is actually part of an infinite family (and considerably simplifies the proof).

A second family of analytical obstructions

Theorem (D. and Van Schaftingen (2024))

For every $n \in \mathbb{N}_*$, if $\dim \mathcal{M} > 4n - 1$, then

$$H_W^{1,4n-1}(\mathcal{M}; \mathbb{S}^{2n}) \subsetneq W^{1,4n-1}(\mathcal{M}; \mathbb{S}^{2n}).$$

The key ingredient is a periodic construction using a Whitehead product.

This shows that Bethuel's counterexample is actually part of an infinite family (and considerably simplifies the proof).

Thank you for your attention!